



2021

Underlying Coronary Microvascular Disease May Lead to Underestimation of Global Longitudinal Strain Differences

Follow this and additional works at: <https://www.j-saudi-heart.com/jsha>



Part of the [Cardiology Commons](#)



This work is licensed under a [Creative Commons Attribution-Noncommercial-No Derivative Works 4.0 License](#).

Recommended Citation

Gupta, Neelesh; Elkhazendar, Mohammed; and Gupta, Rajeev (2021) "Underlying Coronary Microvascular Disease May Lead to Underestimation of Global Longitudinal Strain Differences," *Journal of the Saudi Heart Association*: Vol. 33 : Iss. 4 , Article 7.

Available at: <https://doi.org/10.37616/2212-5043.1272>

This Letter to the Editor is brought to you for free and open access by Journal of the Saudi Heart Association. It has been accepted for inclusion in Journal of the Saudi Heart Association by an authorized editor of Journal of the Saudi Heart Association.

Underlying Coronary Microvascular Disease May Lead to Underestimation of Global Longitudinal Strain Differences

Neelesh Gupta ^a, Mohammed Elkhazendar ^{b,*}, Rajeev Gupta ^c

^a Department of Cardiology, Kirk Kerkorian School of Medicine, University of Nevada, Las Vegas, NV, USA

^b Clinical Sciences Department, College of Medicine, University of Sharjah, United Arab Emirates

^c Department of Cardiology, Mediclinic Al Jowhara Hospital, Al Ain, United Arab Emirates

To the Editor:

We read with interest the original article by Srinivasan G, et al. [1]. In the study, the methodology stated that “Participants with normal coronaries and non-obstructive disease were taken as the control group.” The speckle-tracking Echocardiography-derived Global Longitudinal Strain (GLS) is a sensitive modality for obtaining the GLS. The participants presenting with chest pain with normal coronaries on angiography and non-obstructive disease accepted as control may have microvascular coronary artery disease affecting the control GLS values, leading to underestimation of the difference between the disease and control group. Studies have shown that patients with coronary microvascular disease are known to have a significant lowering of coronary reserves, thus affecting patterns of GLS [2,3]. Thereby, this should

be taken into consideration when interpreting the results of this study.

Conflict of interest

None declared.

References

- [1] Giridharan S, Aashish A, Prasath P, Karthikeyan S, Rangasamy S, AmirthaGanesh B, et al. Two-dimensional Speckle -Tracking Echocardiography derived longitudinal strain in patients with Unstable angina and normal left ventricular ejection fraction. *J Saudi Heart Assoc* 2021;33: 138–46. <https://doi.org/10.1016/j.jjcc.2020.02.005>.
- [2] Jovanovic I, Tesic M, Giga V, Dobric M, Boskovic N, Vratonjic J, et al. Impairment of coronary flow velocity reserve and global longitudinal strain in women with cardiac syndrome X and slow coronary flow. *J Cardiol* 2020;76(1):1–8. <https://doi.org/10.1016/j.jjcc.2020.02.005>.
- [3] Venkataraman P, Wright L, Huynh Q, Marwick TH. Independence of coronary artery disease to subclinical left ventricular dysfunction. *Echocardiography* 2020;37(5):678–87. <https://doi.org/10.1111/echo.14657>.

Author contribution: Conception and design of Study: Neelesh Gupta, Mohammed Elkhazendar, Rajeev Gupta. Literature review: Neelesh Gupta, Mohammed Elkhazendar, Rajeev Gupta. Drafting of manuscript: Rajeev Gupta. Revising and editing the manuscript critically for important intellectual contents: Neelesh Gupta, Mohammed Elkhazendar, Rajeev Gupta. Supervision of the research: Rajeev Gupta.



Received 23 July 2021; accepted 7 August 2021.
Available online 15 October 2021

* Corresponding author.

<https://doi.org/10.37616/2212-5043.1272>

2212-5043/© 2021 Saudi Heart Association. This is an open access article under the CC-BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).